

Thermalization of Gluons with Bose-Einstein Condensation

Zhe Xu,¹ Kai Zhou,^{1,2} Pengfei Zhuang,¹ and Carsten Greiner²

¹*Department of Physics, Tsinghua University and Collaborative Innovation Center of Quantum Matter, Beijing 100084, China*

²*Institut für Theoretische Physik, Johann Wolfgang Goethe-Universität Frankfurt, Max-von-Laue-Strasse 1, 60438 Frankfurt am Main, Germany*

We study the thermalization of gluons far from thermal equilibrium in relativistic kinetic theory. The initial distribution of gluons is assumed to resemble that in the early stage of ultrarelativistic heavy ion collisions. Only elastic scatterings in static, nonexpanding gluonic matter are considered. At first we show that the occurrence of condensation in the limit of vanishing particle mass requires a general constraint for the scattering matrix element. Then the thermalization of gluons with Bose-Einstein condensation is demonstrated in a transport calculation. We see a continuously increasing overpopulation of low energy gluons, followed by a decrease to the equilibrium distribution, when the condensation occurs. The times of the completion of the gluon condensation and of the entropy production are calculated. These times scale inversely with the energy density.

A new state of matter composed of quarks and gluons, the quark-gluon plasma (QGP), has been produced in experiments of ultrarelativistic heavy ion collisions at the Relativistic Heavy Ion Collider (RHIC) and at the Large Hadron Collider (LHC) [1–3]. The shear viscosity over the entropy density (η/s) of the QGP, extracted from the flow measurements, is a small number [4], which indicates that the QGP is an almost perfect fluid. However, the initially produced quarks and gluons are far from thermal equilibrium. How these partons thermalize toward QGP within a short time of about 1 fm/c is still an open question, although different approaches [5, 6] have been developed to study this issue.

Recently, a new idea of the formation of Bose-Einstein condensates in ultrarelativistic heavy ion collisions [7] raises again the interest in thermalization of quarks and gluons produced in such reactions. The condition for a potential condensation is an overoccupied initial distribution of gluons, when compared with the thermal Bose-Einstein distribution with a temperature obtained by assuming a sudden equilibration with energy and gluon number conservation. Quantitative studies [8, 9] have been performed within the kinetic transport theory, where an overoccupied initial condition is given by the color glass condensate formed in high energy heavy ion collisions. All these works were only able to address the evolution of the gluon system till the onset of Bose-Einstein condensation. Whether or not a gluon condensate appears from kinetic statistical studies is an important question. In this Letter we evolve the gluon system beyond the onset toward the full thermalization with a complete Bose-Einstein condensation. It is an important step forward toward understanding the parton thermalization in ultrarelativistic heavy ion experiments. Our work demonstrates an example of Bose-Einstein condensation and thermalization in a relativistic system being far from thermal equilibrium and with high energy density, which also provides new insights for other field of physics.

In this work, the space-time evolution of gluons with the Bose-Einstein condensation is described in a kinetic transport approach BAMPs (Boltzmann approach of

multiparton scatterings) [6] that solves the Boltzmann equation for gluons. We consider a static, nonexpanding gluon system that is initially out of equilibrium and assume gluon number conservation by including only gluonic elastic scatterings in the collision term of the Boltzmann equation. Once the initial distribution of gluons is given, the gluon system will finally evolve to the thermal equilibrium distribution

$$f_{eq}(\mathbf{p}) = \frac{1}{e^{E/T_{BE}} - 1} + (2\pi)^3 n_c^{eq} \delta^{(3)}(\mathbf{p}), \quad (1)$$

which is the solution of the Boltzmann equation in the long time limit. The temperature T_{BE} and the density of the gluon condensate at equilibrium n_c^{eq} are obtained by using the assumed gluon number and energy conservation. We will show that our numerical solution agrees with Eq. (1), and will determine the time scale of the full thermalization for various initial conditions from weak to strong overoccupation.

We note that although particle number changing processes may prevent the Bose-Einstein condensation, studies within field theories including particle production and annihilation processes showed the emergence of a condensate for an intermediate time window [10]. Also, results from kinetic transport calculations indicate an acceleration of the onset of gluon condensation due to gluon bremsstrahlung processes [11].

The formation of a Bose-Einstein condensate is the consequence of quantum statistics of bosons and has been observed in experiments using ultracold atoms [12]. The Boltzmann equation including Bose statistics reads

$$\begin{aligned} \left(\frac{\partial}{\partial t} + \frac{\mathbf{p}_i}{E_i} \frac{\partial}{\partial \mathbf{r}} \right) f_i &= \frac{1}{2E_1} \int d\Gamma_2 \frac{1}{2} \int d\Gamma_3 d\Gamma_4 |\mathcal{M}_{34 \rightarrow 12}|^2 \\ &\times [f_3 f_4 (1 + f_1)(1 + f_2) - f_1 f_2 (1 + f_3)(1 + f_4)] (2\pi)^4 \\ &\times \delta^{(4)}(p_3 + p_4 - p_1 - p_2), \end{aligned} \quad (2)$$

where $f_i = f_i(\mathbf{r}, \mathbf{p}_i, t)$ and $d\Gamma_i = d^3p_i/(2E_i)/(2\pi)^3$, $i = 1, 2, 3, 4$. Binary collisions $34 \rightarrow 12$ and $12 \rightarrow 34$ are determined by the collision kernel $|\mathcal{M}_{34 \rightarrow 12}|^2$ and $|\mathcal{M}_{12 \rightarrow 34}|^2$, which are equal. $(1 + f_1)(1 + f_2)$ and

$(1 + f_3)(1 + f_4)$ are the Bose factors, with which the distribution (1) is a solution of Eq. (2).

We decompose the distribution function f into two parts $f = f^{gas} + f^{cond}$, where f^{gas} denotes the distribution of gas (noncondensate) particles, while $f^{cond} = (2\pi)^3 n_c \delta^{(3)}(\mathbf{p})$ denotes the distribution of the condensate particles with zero momentum. n_c is the density of the condensate particles, which grows with time during the condensation. We obtain easily the equation describing the condensation of zero momentum particles, when we replace f_1 and $1 + f_1$ in Eq. (2) by $f_1^{cond}(\mathbf{r}, \mathbf{p}_1, t)$. Integrating this equation over $\mathbf{p}_1$ gives the time derivative of n_c . After a lengthy integration with the help of the delta functions $\delta^{(3)}(\mathbf{p})\delta^{(4)}(p_3 + p_4 - p_1 - p_2)$ we have (as that derived in [13])

$$\frac{\partial n_c}{\partial t} = \frac{n_c}{64\pi^3} \int dE_3 dE_4 [f_3 f_4 - f_2(1 + f_3 + f_4)] \times E \left[\frac{|\mathcal{M}_{34 \rightarrow 12}|^2}{s} \right]_{s=2mE}. \quad (3)$$

The two terms on the right-hand side correspond to kinetic processes for the condensation and the evaporation, respectively. In the derivation we have assumed that f only depends on the absolute value of $\mathbf{p}$. The term of the spacial derivative of n_c drops out, since we consider a homogeneous gluon matter. m denotes the particle mass at rest. $E = E_3 + E_4$ is the total energy in the collision, while $p = |\mathbf{p}_3 + \mathbf{p}_4|$ is the total momentum. $s = E^2 - p^2$ is the invariant mass. The energy-momentum conservation leads to $E = m + E_2$ and $p = p_2$, where particle 1 denotes the condensate particle with zero momentum. From $E_2^2 = p_2^2 + m^2$ we obtain $s = 2mE$, which is the kinematic constraint for condensation processes.

Equation (3) indicates a general condition for the occurrence of the Bose-Einstein condensation: $|\mathcal{M}|^2/s$ must be finite. In particular, this condition should hold for vanishing rest mass of particles $m = 0$, which is the case we are considering in this work. The elastic scatterings of massless gluons are described in leading order of perturbative QCD. We use

$$|\mathcal{M}_{gg \rightarrow gg}|^2 \approx 144\pi^2 \alpha_s^2 \frac{s^2}{t(t - m_D^2)} \quad (4)$$

as the collision kernel, which has been calculated by using the standard Hard-Thermal-Loop (HTL) treatment according to a sum rule satisfied by the HTL gluon propagator [14, 15]. m_D is the Debye screening mass. In a process $34 \rightarrow 12$, where “1” denotes a condensate gluon with zero momentum and zero mass, gluon 2, 3 and 4 must be colinear. Thus, the Mandelstam variables s and t are zero and s/t goes to -1 . We see that $|\mathcal{M}_{gg \rightarrow gg}|^2/s|_{s=0} \sim \alpha_s^2/m_D^2$ is finite. On the contrary, the often used kernel [16] $|\tilde{\mathcal{M}}|^2 \sim s^2/(t - m_D^2)^2$ gives $|\tilde{\mathcal{M}}|^2/s|_{s=0} = 0$, which indicates that the gluon condensation will not happen, although the gluon system has evolved to the onset.

We note that for scatterings involving noncondensate particles only, i.e., $s > 0$, the total cross section $\sigma_{gg \rightarrow gg}$ is logarithmically divergent due to Eq. (4). This is regularized by an upper cutoff of t , since scatterings with t approaching to zero do not contribute to thermalization.

We have derived Eq. (3) in order to prove the justification of the chosen collision kernel Eq. (4). The time evolution of the condensation and thermalization is achieved by solving the Boltzmann equation (2) numerically. To do it we modify the currently used partonic transport model BAMPs [6] to include the Bose factors. A new scheme [17] is employed to simulate collisions stochastically between test particles. Instead of a collision probability we define a differential collision probability

$$\frac{dP_{22}}{d\Omega} = \frac{v_{rel}}{N_{test}} \frac{d\sigma_{22}}{d\Omega} (1 + f_3)(1 + f_4) \frac{\Delta t}{\Delta V}. \quad (5)$$

$v_{rel} = s/(2E_1 E_2)$ is the relative velocity of the incoming particles, N_{test} the number of test particles per real particle, ΔV the volume element around the collision point, Δt the time step in the calculation, and $d\sigma_{gg \rightarrow gg}/d\Omega$ the differential cross section of elastic gluonic scatterings. At first a solid angle $\tilde{\Omega}$ is sampled according to $d\sigma_{22}/d\Omega$ for each of the particle pairs considered. In case of the occurrence of a collision the momenta $\mathbf{p}_3, \mathbf{p}_4$ of the outgoing particles are thus determined. Kinematic constraints ensure exact energy-momentum conservation in each of the collisions. We also obtain f_3 and f_4 from the extracted f at $\mathbf{p}_3$ and $\mathbf{p}_4$, respectively. Second, a random number between zero and the value of a normalized reference function $dF/d\Omega$ at $\tilde{\Omega}$ will decide whether a collision will occur: if this random number is smaller than the differential collision probability $dP_{22}/d\Omega$ at $\tilde{\Omega}$, a collision occurs; otherwise not. In processes $12 \rightarrow 3c$ where a condensate gluon c is produced, the two incoming gluons should have parallel momenta, which is impossible in numerical calculations. We define a cutoff energy ϵ . Gluons with energy smaller than ϵ are regarded as condensate gluons. Since the production rate of the condensate particles should not depend on the particular numerical implementation, ϵ has to be chosen sufficiently small to avoid numerical artifacts. The details and numerical confirmation of the new scheme will be shown in a forthcoming paper [18].

The momentum distribution of initial gluons is assumed to be

$$f_{init}(\mathbf{p}) = f_0 \theta(Q_s - |\mathbf{p}|), \quad (6)$$

which resembles that in the early stage of ultrarelativistic heavy ion collisions [7]. The initial momentum distribution has been simplified to be isotropic, although the new method introduced [18] can be applied for anisotropic momentum distributions. From (6) we obtain the temperature at equilibrium and the condensate fraction of the total density

$$T_{BE} = (15f_0/4)^{1/4} Q_s/\pi, \quad n_c^{eq}/n = 1 - 19.43\pi^{-3} f_0^{-1/4}. \quad (7)$$

There is a critical value of $f_0^c = 0.154$, below which n_c^{eq}/n becomes negative and no Bose-Einstein condensation occurs. We set $Q_s = 1$ GeV and vary f_0 from 0.4 to 2.0 to change the number and energy density. Gluons are homogeneously distributed in a box, for which the side length is set to be 3 fm. A periodic boundary condition is taken. The box is divided by cubic cells with equal volume ΔV . The side length of a cell is 0.125 fm. Large values of N_{test} are used to enhance statistics and to reduce numerical uncertainties in the extraction of f . For scatterings a constant coupling $\alpha_s = 0.3$ is used and the screening mass is evaluated at each of the time steps by [16]

$$m_D^2 = 16\pi N_c \alpha_s \int \frac{d^3 p}{(2\pi)^3} \frac{1}{p} f^{gas}. \quad (8)$$

Figure 1 shows the energy distributions of noncondensate gluons at various times. $f_0 = 2.0$ is chosen in the




FIG. 1: (color online) The energy distributions of noncondensate gluons at various times. $f_0 = 2$ is chosen.

example. From the upper panel of Fig. 1 we observe a continuous increase at low energy and at energy higher than Q_s , whereas between some intermediate energy and Q_s the distribution decreases continuously. This indicates that particle and energy are flowing from the scale at Q_s toward both lower and higher energies at early times of the thermalization process. The distributions shown in Fig. 1 are calculated by counting test particles

in equal energy intervals of $\Delta p = 0.005$ GeV and dividing these particle numbers by $N_{test} V p^2 \Delta p / (2\pi^2)$, where V is the volume of the box and p is the middle position of the energy intervals. The distributions are plotted at each momenta p . The open symbols in Fig. 1 are first ten values of these distributions. The particle number in intervals at low energies is smaller and thus has a larger fluctuation compared to those at intermediate energies. For the present setups it is difficult to make an unambiguous parametrization for the distributions at low energies. One needs a larger value of N_{test} for obtaining higher accuracy. Nevertheless, we fit the distributions at low energies by Bose-Einstein distributions with different effective temperatures T_{eff} and chemical potentials μ_{eff} . We find that μ_{eff} increases from a negative value to zero, which indicates the onset of the Bose-Einstein condensation [8, 9]. When μ_{eff} is close to zero, its value will fluctuate around zero due to the numerical uncertainty. Once μ_{eff} becomes positive, we define this moment as the start time of the gluon condensation τ_c . At this start time n_c probably has a nonzero value, because we have made the approximation to regard particles with energy smaller than ϵ as condensate gluons. In the calculations we set $\epsilon = 0.0025$ GeV, which is small enough to reduce numerical errors. We see later in Fig. 2 that n_c at long time agrees perfectly with the analytical equilibrium value n_c^{eq} .

We note that both τ_c and n_c at τ_c should be determined from a theory describing the phase transition to a Bose-Einstein condensate [10], which is beyond the scope of the Boltzmann approach. However, we will see these values will not affect the dynamic behavior of the growth of the condensate appreciably. In the calculation for $f_0 = 2.0$ the condensation starts at $\tau_c = 0.376$ fm/c. The energy distribution at τ_c is far from thermal equilibrium. This particular feature differs from the production of Bose-Einstein condensates in experiments using ultra-cold atoms [12], where the system is close to thermal equilibrium. From the upper panel of Fig. 1 we also see that until 0.5 fm/c the condensation is too weak to reduce the increasing overpopulation at low energies. T_{eff} increases to 10 GeV at 0.5 fm/c, which is much larger than the equilibrium temperature, $T_{BE} = 0.53$ GeV.

After 0.5 fm/c the overpopulation at low energies begins to disappear due to the growth of the gluon condensate, as clearly seen in the lower panel of Fig. 1. We find that the distribution at low energies has a profile $f \sim 1/p$ from τ_c . The freed energy during the condensation flows to particles with high energies, where the occupation number increases continuously. The energy distribution relaxes to the equilibrium form. We find a perfect agreement with the equilibrium distribution at 3.4 fm/c, which indicates the completion of the gluon condensation, see Fig. 2. We notice that the distribution at 1.5 fm/c is already very close to the equilibrium one.

The growth of the gluon condensate in time is shown in the upper panel of Fig. 2. At early times of the condensation the gluon condensate grows exponentially, which can




FIG. 2: (color online) Time evolution of the density of the gluon condensate (upper panel) and the entropy production (lower panel). Calculations are performed for various f_0 .

qualitatively be understood by Eq. (3). At these times the production processes are dominant. When the evaporation processes begin to balance the production processes, the growth of the gluon condensate slows down, and then has a relaxation form.

Numerical calculations are performed for three different f_0 . Larger f_0 leads to larger particle and energy densities and larger fraction of the density of the gluon condensate to the total gluon density, see Eq. (7). To study the scaling behavior, the density of the gluon condensate is divided by its final value at equilibrium, while the time is multiplied by f_0 . We see an approximate scaling, which shows that *the larger the density (larger f_0), the faster is the completion of the gluon condensation, and the faster is the thermal equilibration*. The scaling behavior of the starting time of the gluon condensation τ_c can be understood by approximating the collision term in the Boltzmann equation (2), $f_3 f_4 (1+f_1)(1+f_2) - f_1 f_2 (1+f_3)(1+f_4) \approx f_3 f_4 (f_1 + f_2) - f_1 f_2 (f_3 + f_4)$. It is a good approximation for colliding particles with small energy, where f is large. If f at low energies is proportional to f_0 , $f \sim f_0 f_N(p, t)$, which seems reasonable at least at early times due to the given initial condition (6), and if this relation holds long for large f_0 , then the right-hand side of (2) scales with f_0^2 by noting that $|\mathcal{M}_{gg \rightarrow gg}|^2 \sim 1/m_D^2$

and m_D^2 is proportional to f_0 . The Boltzmann equation (2) can be reduced to an equation for $f_N(p, t')$ with $t' = f_0 t$. Since $f_N(p, t') \approx T_{eff}(t')/f_0/[p - \mu_{eff}(t')]$ for low energy, μ_{eff} depends on t' , which explains the scaling behavior of τ_c , when μ_{eff} approaches zero with t' . In addition, T_{eff} is proportional to f_0 at τ_c , which is also confirmed from our numerical results. After τ_c the scaling of n_c can be understood analogously by using $f_3 f_4 - f_2(1+f_3+f_4) \approx f_3 f_4 - f_2(f_3+f_4)$ in Eq. (3), if the relation $f \sim f_0 f_N$ still holds. At late times when n_c relaxes to its final value, f relaxes to the Bose-Einstein distribution and is not proportional to f_0 even at low energies. This violates slightly the time scaling of n_c , which is seen in Fig. 2 at late times. On the contrary, the differences of $n_c(t')$ shortly after τ_c are due to numerical uncertainties in τ_c as well as n_c at τ_c .

The lower panel of Fig. 2 shows the entropy production $s(t) - s(t_0 = 0)$ divided by its value at equilibrium. We see a two step production separated by τ_c . This shows that the formation of the gluon condensate is essential for the time scale of thermalization. From Fig. 2 we also see that the completion of the condensation is later than that of entropy production. The full thermalization is reached at $f_0 t \approx 3$ fm/c. In addition, the entropy production does not scale with $f_0 t$ before τ_c , because the momentum distribution at low energies has a negligible contribution to the entropy and the argument for τ_c scaling does not apply to the early entropy production. Instead, we find that the early entropy production is independent of f_0 . The reason lies in the almost same collision rate at the early times, because $\sigma_{gg \rightarrow gg} \sim 1/m_D^2$ and m_D^2 is roughly proportional to f_0 .

In this Letter we demonstrated for the first time a complete Bose-Einstein condensation of gluons within kinetic theory. Our calculations showed that the times for the occurrence of the condensation, the completion of the condensation, and the full thermalization scale inversely with the energy density. For instance, for energy density 53 GeV/fm^3 ($f_0 = 2$) the time of the full thermalization is about 1.5 fm/c, which is consistent with the parametric result $t_{th} \sim Q_s^{-1} \alpha_s^{-7/4}$ found in [7]. We also find that the earlier the condensation occurs, the earlier does the second step of the entropy production start. All these are important new findings to get further forward with the understanding of thermalization for systems far from thermal equilibrium.

As discussed in [19–21], pions, photons, and dileptons produced in ultrarelativistic heavy ion collisions might provide measurable hints for the formation of a gluon condensate. The experimental significance also depends on how long the gluon condensate lives after its formation. In our present study a gluon condensate exits, because we assumed the gluon number conservation and considered a static, nonexpanding system. The inclusion of quark-antiquark annihilations [22], or/and gluon bremsstrahlung processes, and the back reactions [11, 23] will violate the gluon number conservation, which will reduce the overoccupation of gluons at small momenta.

Whether a gluon condensation will still occur, is an interesting question. If the condensation in fact happens, the gluon condensate is only an intermediate state and may disappear in the later evolution. Moreover, one has to consider the expansion of quark gluon matter, in order to obtain more realistic results. All these are subjects of future works.

Z.X. thanks X.G. Huang, J. Liao, V. Greco, and L. McLerran for helpful discussions. This work was financially supported by the MOST, the NSFC under Grants No. 2014CB845400, No. 11275103, and No. 11335005. The BAMPS simulations were performed at Tsinghua National Laboratory for Information Science and Technology.

-
- [1] J. Adams *et al.* (STAR Collaboration), Nucl. Phys. A **757**, 102 (2005) [nucl-ex/0501009].
 - [2] K. Adcox *et al.* (PHENIX Collaboration), Nucl. Phys. A **757**, 184 (2005) [nucl-ex/0410003].
 - [3] K. Aamodt *et al.* (ALICE Collaboration), Phys. Rev. Lett. **105**, 252302 (2010) [arXiv:1011.3914 [nucl-ex]].
 - [4] B. Schenke, S. Jeon, and C. Gale, Phys. Lett. B **702**, 59 (2011) [arXiv:1102.0575 [hep-ph]]; Phys. Rev. C **85**, 024901 (2012) [arXiv:1109.6289 [hep-ph]]; H. Song, S. A. Bass, U. Heinz, T. Hirano, and C. Shen, Phys. Rev. Lett. **106**, 192301 (2011) [Erratum-ibid. **109**, 139904 (2012)] [arXiv:1011.2783 [nucl-th]]; P. Romatschke, Int. J. Mod. Phys. E **19**, 1 (2010) [arXiv:0902.3663 [hep-ph]].
 - [5] X. G. Huang and J. Liao, Int. J. Mod. Phys. E **23**, 1430003 (2014) [arXiv:1402.5578 [nucl-th]]; M. Strickland, arXiv:1312.2285 [hep-ph]; F. Gelis, Int. J. Mod. Phys. A **28**, 1330001 (2013) [arXiv:1211.3327 [hep-ph]]; J. Berges, J. P. Blaizot, and F. Gelis, J. Phys. G **39**, 085115 (2012) [arXiv:1203.2042 [hep-ph]]; P. B. Arnold, Int. J. Mod. Phys. E **16**, 2555 (2007) [arXiv:0708.0812 [hep-ph]]; D. Bodeker and K. Rummukainen, JHEP **0707**, 022 (2007) [arXiv:0705.0180 [hep-ph]].
 - [6] Z. Xu and C. Greiner, Phys. Rev. C **71**, 064901 (2005) [arXiv:hep-ph/0406278]; **76**, 024911 (2007) [arXiv:hep-ph/0703233].
 - [7] J. P. Blaizot, F. Gelis, J. F. Liao, L. McLerran, and R. Venugopalan, Nucl. Phys. A **873**, 68 (2012) [arXiv:1107.5296 [hep-ph]].
 - [8] J. P. Blaizot, J. Liao, and L. McLerran, Nucl. Phys. A **920**, 58 (2013) [arXiv:1305.2119 [hep-ph]].
 - [9] F. Scardina, D. Perricone, S. Plumari, M. Ruggieri, and V. Greco, Phys. Rev. C **90**, no. 5, 054904 (2014) [arXiv:1408.1313 [nucl-th]].
 - [10] T. Epelbaum and F. Gelis, Nucl. Phys. A **872**, 210 (2011) [arXiv:1107.0668 [hep-ph]]; J. Berges and D. Sexty, Phys. Rev. Lett. **108**, 161601 (2012) [arXiv:1201.0687 [hep-ph]]; A. Kurkela and G. D. Moore, Phys. Rev. D **86**, 056008 (2012) [arXiv:1207.1663 [hep-ph]]; J. Berges, K. Boguslavski, S. Schlichting, and R. Venugopalan, Phys. Rev. Lett. **114**, no. 6, 061601 (2015) [arXiv:1408.1670 [hep-ph]].
 - [11] X. G. Huang and J. Liao, arXiv:1303.7214 [nucl-th].
 - [12] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, Science **269**, 198 (1995); K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, Phys. Rev. Lett. **75**, 3969 (1995).
 - [13] D. V. Semikoz and I. I. Tkachev, Phys. Rev. Lett. **74**, 3093 (1995) [hep-ph/9409202].
 - [14] P. Aurenche, F. Gelis, and H. Zaraket, JHEP **0205**, 043 (2002) [hep-ph/0204146].
 - [15] A. Kurkela and G. D. Moore, JHEP **1112**, 044 (2011) [arXiv:1107.5050 [hep-ph]].
 - [16] S. M. H. Wong, Phys. Rev. C **54**, 2588 (1996) [hep-ph/9609287].
 - [17] B. Jackson and E. Zaremba, Phys. Rev. A **66**, 033606 (2002).
 - [18] K. Zhou, Z. Xu, P. Zhuang, and C. Greiner, in preparation.
 - [19] D. N. Voskresensky, J. Exp. Theor. Phys. **78**, 793 (1994); Phys. Atom. Nucl. **59**, 2015 (1996); E. E. Kolomeitsev, B. Kampfer, and D. N. Voskresensky, Acta Phys. Polon. B **27**, 3263 (1996).
 - [20] M. Chiu, T. K. Hemmick, V. Khachatryan, A. Leonidov, J. Liao, and L. McLerran, Nucl. Phys. A **900**, 16 (2013) [arXiv:1202.3679 [nucl-th]].
 - [21] L. McLerran and B. Schenke, Nucl. Phys. A **929**, 71 (2014) [arXiv:1403.7462 [hep-ph]].
 - [22] J. P. Blaizot, B. Wu, and L. Yan, Nucl. Phys. A **930**, 139 (2014) [arXiv:1402.5049 [hep-ph]].
 - [23] M. C. Abreu York, A. Kurkela, E. Lu, and G. D. Moore, Phys. Rev. D **89**, no. 7, 074036 (2014) [arXiv:1401.3751 [hep-ph]]; A. Kurkela and E. Lu, Phys. Rev. Lett. **113**, no. 18, 182301 (2014) [arXiv:1405.6318 [hep-ph]].